



NDC 0904-6349-46

REGULAR
STRENGTH**RANITIDINE**
75**Ranitidine Tablets USP, 75 mg**
Acid Reducer**Prevents and Relieves**
Heartburn Associated with
Acid Indigestion and Sour Stomach**30 Tablets****IMPORTANT:** Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.**Active ingredient (in each tablet)** **Purpose**

Ranitidine 75 mg

(as ranitidine hydrochloride USP, 84 mg).....Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers.**Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.

■ heartburn with **lightheadedness, sweating or dizziness**

■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

■ frequent **chest pain**

■ frequent wheezing, particularly with heartburn

■ unexplained weight loss

■ nausea or vomiting

■ stomach pain

Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).**Directions**

■ adults and children 12 years and over:

■ to **relieve** symptoms, swallow 1 tablet with a glass of water■ to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn.

■ can be used up to twice daily (do not take more than 2 tablets in 24 hours)

■ do not chew tablet

■ children under 12 years: ask a doctor

Other information■ **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**

■ store at 20° - 25° C (68° - 77° F)

■ avoid excessive heat or humidity

■ this product is sodium and sugar free

Inactive ingredients

See carton for a complete list of inactive ingredients

Questions? call 1-800-406-7984Distributed By **MAJOR® PHARMACEUTICALS**

17177 N Laurel Park Drive, Suite 233, Livonia, MI 48152

M-59 Re-Order No. 016074

Rev. 10/16

MADE IN INDIA

Batch No.

Expiration Date:



5138658

Non Varnish Area