

DG7984R1

No print/No varnish
Lot & Exp date

*This product is not manufactured or distributed by Meilei Consumer Healthcare, owner of the registered trademark Tylenol® SINUS + HEADACHE. 50944 REV03/6F46608

Distributed by MAJOR* PHARMACEUTICALS
31778 Enterprise Drive, Livonia, MI 48150 USA M-17
Rev. 06/16 Re-order No. 100491

NDC 0904-5-783-24
*Compare to the active ingredients
in Tylenol® SINUS + HEADACHE

MAJOR®

MAXIMUM STRENGTH Māpap® Sinus CONGESTION AND PAIN CAPLETS

Acetaminophen - Pain reliever / Fever reducer
Phenylephrine HCl - Nasal decongestant

Relieves Headache Pain,
Sinus Pressure & Nasal Congestion



24 CAPLETS with Cool
Blast Flavor

NON-DROWSY
DAYTIME

Drug Facts
KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT
INFORMATION
Active ingredients
(In each caplet)
Acetaminophen 325 mg.....Pain reliever/fever reducer
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses
Temporarily relieves these symptoms associated
with hay fever or other respiratory allergies, and the
common cold:
■ sinus congestion and pressure
■ headache
■ minor aches and pains
■ temporarily relieves these symptoms associated
with hay fever or other respiratory allergies, and the
common cold:
■ sinus congestion and pressure

Drug Facts (continued)
Inactive ingredients
corn starch,
croscopolone, D&C yellow #10 aluminum lake, FD&C
blue #1 aluminum lake, FD&C red #40 aluminum lake,
flavor, magnesium stearate, microcrystalline cellulose,
polyethylene glycol, polyvinyl alcohol, povidone, *

Drug Facts (continued)
Inactive ingredients
silica gel, sodium starch glycolate, stearic acid,
sucralose, talc, titanium dioxide

Questions or comments? (800) 616-2471

Drug Facts (continued)
Warnings
Liver warning: This product contains acetaminophen.
Severe liver damage may occur if you take:
■ more than 4,000 mg in 24 hours, which is the
maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this
product
Allergy alert: Acetaminophen may cause severe skin
reactions. Symptoms may include:
■ hives ■ rash ■ skin reddening
If a skin reaction occurs, stop use and seek medical
help right away.
Do not use
■ with any other drug containing acetaminophen
(prescription or nonprescription). If you are not sure
whether a drug contains acetaminophen, ask a
doctor or pharmacist.
■ if you are now taking a prescription monoamine
oxidase inhibitor (MAOI) (certain drugs for
depression, psychiatric or emotional conditions, or
Parkinson's disease), or for 2 weeks after stopping
the MAOI drug. If you do not know if you
prescription drug contains an MAOI, ask a doctor or
pharmacist before taking this product.
■ if you have ever had an allergic reaction to this
product or any of its ingredients
Ask a doctor before use if you have ■ liver disease
■ difficulty in urination due to enlargement of the
prostatic gland ■ heart disease ■ diabetes

Drug Facts (continued)
■ high blood pressure
■ thyroid disease
Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin.
When using this product do not exceed
recommended dosage.
Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ pain or nasal congestion gets worse or lasts more
than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health
professional before use.
Keep out of reach of children. In case of accidental
overdose, get medical help or contact a Poison
Control Center (1-800-222-1222) right away. Prompt
medical attention is critical for adults as well as for
children even if you do not notice any signs or
symptoms.
Directions ■ do not take more than directed
■ adults and children 12 years and over
■ take 2 caplets every 4 hours
■ swallow whole – do not crush, chew, or dissolve
■ do not take more than 10 caplets in 24 hours
■ children under 12 years: ask a doctor
Other information ■ TAMPER EVIDENT: DO
NOT USE IF OUTER PACKAGE IS OPENED OR
BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between
15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING

B-1212-466C-08SC
REV03/6F46608