

No print/No varnish
Lot & Exp date

3



6

Rev. 06/19

A-17

Re-order No. 37077

TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

B-12-227-12-RU
ORG031822712

Drug Facts (continued)

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for
- diabetes
- arthritis
- gout

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
 - you experience any of the following signs of stomach bleeding: ■ vomit blood ■ feel faint
 - have bloody or black stools
 - have stomach pain that does not get better
 - ringing in the ears or a loss of hearing occurs
 - new symptoms occur
 - pain gets worse or lasts more than 10 days
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless explicitly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-800-222-2222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

cellulose, colloidal anhydrous silica, corn starch, D&G yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polyacrylate, polyethylene glycol, polyvinyl acetate phthalate, polyethylene glycol, stearic wax, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-645-2158

This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Ecotrin®.

5084 ORG031822712

Distributed by: **RUGBY® LABORATORIES**
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152

www.rugbylaboratories.com

Rugby®

Regular Strength
Aspirin 325 mg
Pain Reliever (NSAID)

Aspirin Regimen

100 Enteric Coated Tablets

Compare to the active ingredient in Ecotrin®†

Talk to your doctor or other
healthcare provider before using
this product for your heart.

Safety Coated

Actual Size



NDC 0536-1232-01

Rugby®

Regular Strength
Aspirin
325 mg
Pain Reliever (NSAID)

Aspirin Regimen

100 Enteric Coated Tablets

Drug Facts

Active ingredient
(in each tablet)

Aspirin 325 mg (NSAID) Pain reliever
nonsteroidal anti-inflammatory drug

Uses For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- hives
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product

SEEK OTHER PACKAGE FOR
COMPLETE PRODUCT INFORMATION

Purpose

DG97A