

## Drug Facts (continued)

### Directions

- take every 4 to 6 hours ■ do not take more than 6 doses in 24 hours
- use only with enclosed dosing cup. Do not use with any other device

|                                       |  |
|---------------------------------------|--|
| adults and children 12 years and over | 10-20mL (25 mg to 50 mg)               |
| children 6 to under 12 years          | 5-10mL (12.5 mg to 25 mg)              |
| children 2 to 5 years of age          | do not use unless directed by a doctor |
| children under 2 years of age         | do not use                             |

### Other information

- each (5 mL) contains: **sodium 7 mg**
- **TAMPER-EVIDENT:** Do not use this product if inner foil seal over the mouth of the bottle is cut, torn, broken or missing
- store at room temperature 20°-25°C (68°-77°F)

**Inactive ingredients** artificial cherry flavor, citric acid, D&C Red #33, FD&C Red #40, glycerin, polysorbate 20, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol solution

**Questions or comments?** (800) 616-2471

Distributed by:  
**MAJOR® PHARMACEUTICALS**  
31778 Enterprise Drive  
Livonia, MI 48150, USA

M-64 Rev. 02/16  
Re-order No. 235192



Lot / Exp.:

\* This product is not distributed or manufactured by Pfizer Consumer HealthCare, Pfizer, Inc. distributor of Benadryl® Allergy Liquid. Benadryl® is a registered trademark of Warner-Lambert Company L.L.C.

**MAJOR®** NDC 0904-1228-00  
TAMPER-EVIDENT™  
**Banophen**  
**ALLERGY**

Antihistamine  
12.5 mg Diphenhydramine HCl USP

ALCOHOL FREE

FOR THE TEMPORARY  
RELIEF OF:

Upper Respiratory Allergies  
Hay Fever  
The Common Cold

Compare to the active ingredient  
of Benadryl® Allergy Liquid\*

4 FL. OZ. (118 mL)

## Drug Facts

**Active ingredient (in each 5 mL)** ..... **Purpose**  
Diphenhydramine HCl USP 12.5 mg ..... Antihistamine

**Uses** temporarily relieves these symptoms of hay fever or other upper respiratory allergies ■ runny nose ■ sneezing  
■ itchy, watery eyes ■ itching of the nose or throat

### Warnings

**Do not use** ■ to make a child sleepy ■ with any other product containing diphenhydramine, including one applied topically.

**Ask a doctor before use if you have**

- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers.

**When using this product** ■ marked drowsiness may occur  
■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**PEEL HERE** ▼