



NDC 0536-1232-01

Regular Strength

Aspirin 325 mg

Pain Reliever (NSAID)

Aspirin Regimen
Safety Coated

100 Enteric
Coated Tablets



Distributed by: **RUGBY® LABORATORIES**
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152 www.rugbylaboratories.com

Rev. 06/19 R-17 Re-order No. 370717 50844 ORG031822712

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient (in each tablet)

Purpose

Aspirin 325 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

Warnings See carton for complete warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling ■ hives ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor

■ children under 12 years: do not use unless directed by a doctor

Other information

■ **TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL**

■ **UNDER CAP IS BROKEN OR MISSING**

■ store at 25°C (77°F); excursion's permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package