

SAVE CARTON FOR COMPLETE DRUG FACTS

Drug Facts

Active ingredient
(in each 0.3 mL)
Simethicone 20 mg Antigas

Use
relieves the symptoms referred to as gas

Warnings
Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Directions

- shake well before using
- all dosages may be repeated as needed, after meals and at bedtime, or as directed by a physician
- do not exceed 12 doses per day
- fill enclosed dropper to recommended dosage level and dispense liquid slowly into baby's mouth, toward the inner cheek
- dosage can also be mixed with 1 oz. of cool water, infant formula or other suitable liquids
- clean dropper well after each use and recap bottle with original cap

Age (years)	Weight (lbs)	Dose
infants (under 2)	under 24	0.3 mL
children (2 and over)	24 and over	0.6 mL

Other information

- **TAMPER-EVIDENT:** Do not use if foil seal under the cap is broken or missing.
- store at room temperature
- do not freeze
- see bottom panel for lot number and expiration date

Drug Facts (continued)

Inactive ingredients
carboxymethylcellulose
sodium, citric acid, flavor,
maltitol, microcrystalline
cellulose, purified water,
sodium benzoate, sodium
citrate, xanthan gum

**Questions or
comments?**
(800) 616-2471

*This product is not manufactured
or distributed by the owner of the
registered trademark MYLICON®.

MAJOR®
INFANTS'
GAS RELIEF
DROPS

NDC 0904-5894-30

MAJOR®
INFANTS'
GAS RELIEF
DROPS

Simethicone Oral Suspension USP



Antigas
Relieves the Symptoms
Referred to as Gas
DYE FREE
Plastic Dropper Enclosed

1 FL OZ
(30 mL)



Compare to
active ingredient
in MYLICON®
Infants Drops*



DOSAGE

Children
2 years
and over

Infants
under
2 years

DROPPER ENCLOSED

Distributed by
MAJOR® PHARMACEUTICALS
17177 N Laurel Park Drive,
Suite 233
Livonia, MI 48152 USA
Rev 12/16 M-99
Re-Order No. 700032
426-26119-4



3 09045 89430 9