

5138655

5138655

Keep the directions, consumer information leaflet and warnings before use.

**Drug Facts (continued)**

**Warnings**  
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

**Do not use**  
■ If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.  
■ with other acid reducers  
■ if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**  
■ had heartburn over 3 months. This may be a sign of a more serious condition  
■ heartburn with **lightheadedness, sweating or dizziness**  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulder; or lightheadedness  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss  
■ nausea or vomiting  
■ stomach pain

**Stop use and ask a doctor if**  
■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days

**Drug Facts (continued)**

**Directions**  
■ adults and children 12 years and over: to **relieve** symptoms, swallow 1 tablet with a glass of water  
■ to **prevent** symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn  
■ can be used up to twice daily (do not take more than 2 tablets in 24 hours)  
■ children under 12 years: ask a doctor

**Other information**  
■ **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.**  
■ store at 20° - 25° C (68° - 77° F)  
■ avoid excessive heat or humidity  
■ this product is sodium and sugar free

**Inactive ingredients**  
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

**Questions?** call 1-800-406-7984

**Drug Facts**

**Active ingredient (in each tablet)**  
Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)

**Purpose**  
Acid reducer

**Uses**  
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages  
■ relieves heartburn associated with acid indigestion and sour stomach

**MAJOR**  
MAXIMUM STRENGTH  
**RANITIDINE 150**  
Ranitidine Tablets USP, 150 mg  
Acid Reducer

NDC 0904-6350-24

**RANITIDINE 150**  
Ranitidine Tablets USP, 150 mg  
Acid Reducer

Prevents and Relieves  
Heartburn Associated with Acid Indigestion and Sour Stomach

**COMPARE TO** the active ingredient of  
**ZANTAC 150**\*\*

Actual Size

**24 Tablets**

\*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Zantac 150®.

Expiration Date:

Batch No.

NON VARNISH



Distributed By  
**MAJOR® PHARMACEUTICALS**  
17177 N Laurel Park Drive, Suite 233  
Livonia, MI 48152  
M-59 Re-Order No. 016076  
**MADE IN INDIA**