NDC 0904-6304-77

## 

## Magnesium Citrate Oral Solution

Effervescent saline laxative Lemon flavored

do not use if tamper evident twist-off cap is Missing, broken or separated from neckring

## 10 FL OZ (296 mL)

Drug Facts	Drug Facts (continued)		s
Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 gSaline laxative	or contact a Poison Control Center right away.		[]CAI
	Store at temperatures between 46° and 86° F (8° and 30° C)		5 S
Use for relief of occasional constipation (irregularity). Generally produces bowel movement in ½ to 6 hours	<b>Directions •</b> drink a full glass (8 ounces) of liquid with each dose <b>•</b> may be taken as a single daily dose or in divided doses		Distributed Bv: MaJOR® PHARMACEUTICALS
Warnings Do not use	adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours	r: MAJOF
If you are on a low sait diet Ask a doctor before use if you have kidney disease im a magnesum or sodium-restricted diet abdominal pain, nausea, or vomiting noticed a sudden change in bowel habits that persists	children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours	buted By
	children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours	Distri
over a period of 2 weeks already used a laxative for a period longer than 1 week	children under 2 years of age	ask a doctor	
Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before as after attend turns. Laureline mou affect hour	discard unused product within 24 hours of opening bottle		
before or after other drugs. Laxatives may affect how other drugs work.	Other information each fl oz contains: magnesium 290 mg each fl oz contains: sodium 45 mg		
Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.			
	Inactive ingredients citric acid, lemon oil, polyethylene glycol, purified water, sodium		∣≡
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help	bicarbonate, sodium saccharin, sucrose		
	Questions or comments? (800) 616-2471		

L0013151SB

Livonia, MI 48 152 Re-Order No. 014432 1 Rev. 10/16

76-M

09046 30477

Μ