


No Print Area
Lot No. & Exp. Date



TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

B-1212-291-13-RU
REV1116A29113

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts	Active ingredient (in each brown tablet) Purpose Ibuprofen USP 200 mg (NSAID) Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug
Uses	temporarily relieves minor aches and pains due to: ■ menstrual cramps ■ toothache ■ backache ■ the common cold ■ headache ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever
Warnings	Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ shock ■ facial swelling ■ blisters ■ rash ■ hives ■ skin reddening ■ asthma (wheezing) If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ take more or for a longer time than directed ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use	■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery
Ask a doctor before use if	■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers

Drug Facts (continued)	Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug
When using this product	■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if	■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ you have stomach pain that does not get better ■ you have symptoms of heart problems or stroke ■ chest pain ■ slurred speech ■ leg swelling ■ trouble breathing ■ weakness in one part or side of body ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear
If pregnant or breast-feeding	ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.
Directions	■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor
Other information	■ store between 20°-25°C (68°-77°F) ■ avoid excessive heat 40°C (104°F) ■ see end flap for expiration date and lot number



NDC 0536-1088-02
Compare to the active ingredient in Advil® Tablets†

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/Fever Reducer
(NSAID)



250 COATED TABLETS



NDC 0536-1088-02
Compare to the active ingredient in Advil® Tablets†

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/Fever Reducer
(NSAID)



250 COATED TABLETS

Drug Facts (continued)

Inactive ingredients
corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? 1-800-645-2158

†This product is not manufactured or distributed by Wyeth Consumer Healthcare, owner of the registered trademark Advil® Tablets.
50844 REV1116A29113
Distributed by: Rugby Laboratories
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152 USA www.rugbylaboratories.com
R-17 Re-order No. 301935 Rev. 09/17

DG982B